

SUBJECT INDEX Volume 64

Adipocytes

Zangani D., Darcy K.M., Masso-Welch P.A., Bellamy E.S., Desole M.S., Ip M.M.
Multiple differentiation pathways of rat mammary stromal cells *in vitro*: acquisition of a fibroblast, adipocyte or endothelial phenotype is dependent on hormonal and extracellular matrix stimulation, 64:91–101

Garcés C., Ruiz-Hidalgo M.J., Bonvini E., Goldstein J., Laborda J.
Adipocyte differentiation is modulated by secreted delta-like (dlk) variants and requires the expression of membrane-associated dlk, 64:103–114

Berberich S.J., Litteral V., Mayo L.D., Tabesh D., Morris D.
mdm-2 gene amplification in 3T3-L1 preadipocytes, 64:205–212

Boone C., Grégoire F., Remacle C.
Various stimulators of the cyclic AMP pathway fail to promote adipose conversion of porcine preadipocytes in primary culture, 64:255–262

Armadillo proteins

Mertens C., Kuhn C., Moll R., Schwetlick I., Franke W.W.
Desmosomal plakophilin 2 as a differentiation marker in normal and malignant tissues, 64:271–290

Schmidt A., Langbein L., Prätzel S., Rode M., Rackwitz H.-R., Franke W.W.
Plakophilin 3 – a novel cell-type-specific desmosomal plaque protein, 64:291–306

arm-repeat proteins

Mertens C., Kuhn C., Moll R., Schwetlick I., Franke W.W.
Desmosomal plakophilin 2 as a differentiation marker in normal and malignant tissues, 64:271–290

Schmidt A., Langbein L., Prätzel S., Rode M., Rackwitz H.-R., Franke W.W.
Plakophilin 3 – a novel cell-type-specific desmosomal plaque protein, 64:291–306

Basal lamina

Tomakidi P., Mirancea N., Fusenig N.E., Herold-Mende C., Bosch F.X., Breitkreutz D.
Defects of basement membrane and hemidesmosome structure correlate with malignant phenotype and stromal interactions in HaCaT-Ras xenografts, 64:263–275

Bile acid synthesis

Stier H., Fahimi H.D., Van Veldhoven P.P., Mannaerts G.P., Völkl A., Baumgart E.
Maturation of peroxisomes in differentiating human hepatoblastoma cells (HepG2): possible involvement of the peroxisome proliferator-activated receptor α (PPAR α), 64:35–66

Body-site specificity of epidermis

Compton C.C., Nadire K.B., Regauer S., Simon M., Warland G., Landry D.B.
Cultured human sole-derived keratinocyte grafts re-express site-specific differentiation after transplantation, 64:45–53

Branched-chain-acyl-CoA oxidase

Stier H., Fahimi H.D., Van Veldhoven P.P., Mannaerts G.P., Völkl A., Baumgart E.
Maturation of peroxisomes in differentiating human hepatoblastoma cells (HepG2): possible involvement of the peroxisome proliferator-activated receptor α (PPAR α), 64:35–66

Bromodeoxyuridine

McBride S., Walsh D., Meleady P., Daly N., Clynes M.
Bromodeoxyuridine induces keratin protein synthesis at a post-transcriptional level in human lung tumour cell lines, 64:185–193

3T3 cells

Garcés C., Ruiz-Hidalgo M.J., Bonvini E., Goldstein J., Laborda J.
Adipocyte differentiation is modulated by secreted delta-like (dlk) variants and requires the expression of membrane-associated dlk, 64:103–114

C3H10T1/2 cells

Denker A.E., Haas A.R., Nicoll S.B., Tuan R.S.
Chondrogenic differentiation of murine C3H10T1/2 multipotential mesenchymal cells: I. Stimulation by bone morphogenetic protein-2 in high-density micromass cultures, 64:67–76

Haas A.R., Tuan R.S.
Chondrogenic differentiation of murine C3H10T1/2 multipotential mesenchymal cells: II. Stimulation by bone morphogenetic protein-2 requires modulation of N-cadherin expression and function, 64:77–89

Cadherins

Haas A.R., Tuan R.S.
Chondrogenic differentiation of murine C3H10T1/2 multipotential mesenchymal cells: II. Stimulation by bone morphogenetic protein-2 requires modulation of N-cadherin expression and function, 64:77–89

cAMP

Boone C., Grégoire F., Remacle C.
Various stimulators of the cyclic AMP pathway fail to promote adipose conversion of porcine preadipocytes in primary culture, 64:255–262

Cardiac muscle

Thorsteinsdóttir S., Roelen B.A.J., Goumans M.-J., Ward-van Oostwaard D., Gaspar A.C., Mummery C.L.
Expression of the $\alpha 6 A$ integrin splice variant in developing mouse embryonic stem cell aggregates and correlation with cardiac muscle differentiation, 64:173–184

Cartilage development

Denker A.E., Haas A.R., Nicoll S.B., Tuan R.S.
Chondrogenic differentiation of murine C3H10T1/2 multipotential mesenchymal cells: I. Stimulation by bone morphogenetic protein-2 in high-density micromass cultures, 64:67–76

Haas A.R., Tuan R.S.
Chondrogenic differentiation of murine C3H10T1/2 multipotential mesenchymal cells: II. Stimulation by bone morphogenetic protein-2 requires modulation of N-cadherin expression and function, 64:77–89

Cdx1

Subramanian V., Meyer B., Evans G.S.
The murine *Cdx1* gene product localises to the proliferative compartment in the developing and regenerating intestinal epithelium, 64:11–18

Cell adhesion

Haas A.R., Tuan R.S.

Chondrogenic differentiation of murine C3H10T1/2 multipotential mesenchymal cells: II. Stimulation by bone morphogenetic protein-2 requires modulation of N-cadherin expression and function, 64:77-89

Cell culture

Stier H., Fahimi H.D., Van Veldhoven P.P., Mannaerts G.P., Völkl A., Baumgart E.

Maturation of peroxisomes in differentiating human hepatoblastoma cells (HepG2): possible involvement of the peroxisome proliferator-activated receptor α (PPAR α), 64:35-66

Cell elongation

Sullivan C.H., Marker P.C., Thorn J.M., Brown J.D.

Reliability of delta-crystallin as a marker for studies of chick lens induction, 64:1-9

Cell fusion

Hirayama E., Nakanishi M., Honda N., Kim J.

Mouse C2 myoblast cells resist HVJ (Sendai virus)-mediated cell fusion in the proliferating stage but become capable of fusion after differentiation, 64:213-223

Chicken embryo

Sullivan C.H., Marker P.C., Thorn J.M., Brown J.D.

Reliability of delta-crystallin as a marker for studies of chick lens induction, 64:1-9

Chondrogenesis

Denker A.E., Haas A.R., Nicoll S.B., Tuan R.S.

Chondrogenic differentiation of murine C3H10T1/2 multipotential mesenchymal cells: I. Stimulation by bone morphogenetic protein-2 in high-density micromass cultures, 64:67-76

Haas A.R., Tuan R.S.

Chondrogenic differentiation of murine C3H10T1/2 multipotential mesenchymal cells: II. Stimulation by bone morphogenetic protein-2 requires modulation of N-cadherin expression and function, 64:77-89

Collagens

Denker A.E., Haas A.R., Nicoll S.B., Tuan R.S.

Chondrogenic differentiation of murine C3H10T1/2 multipotential mesenchymal cells: I. Stimulation by bone morphogenetic protein-2 in high-density micromass cultures, 64:67-76

Haas A.R., Tuan R.S.

Chondrogenic differentiation of murine C3H10T1/2 multipotential mesenchymal cells: II. Stimulation by bone morphogenetic protein-2 requires modulation of N-cadherin expression and function, 64:77-89

Crypts

Subramanian V., Meyer B., Evans G.S.

The murine *Cdx1* gene product localises to the proliferative compartment in the developing and regenerating intestinal epithelium, 64:11-18

Crystallins

Sullivan C.H., Marker P.C., Thorn J.M., Brown J.D.

Reliability of delta-crystallin as a marker for studies of chick lens induction, 64:1-9

Mizuno N., Mochii M., Takahashi T.C., Eguchi G., Okada T.S. Lens regeneration in *Xenopus* is not a mere repeat of lens development, with respect to crystallin gene expression, 64:143-149

Cultured epithelial autograft

Compton C.C., Nadire K.B., Regauer S., Simon M., Warland G., Landry D.B.

Cultured human sole-derived keratinocyte grafts re-express site-specific differentiation after transplantation, 64:45-53

Desmosomes

Mertens C., Kuhn C., Moll R., Schwetlick I., Franke W.W.

Desmosomal plakophilin 2 as a differentiation marker in normal and malignant tissues, 64:271-290

Schmidt A., Langbein L., Prätzel S., Rode M., Rackwitz H.-R., Franke W.W.

Plakophilin 3 - a novel cell-type-specific desmosomal plaque protein, 64:291-306

Development

Mills A.A., Mills M.J., Gardiner D.M.,

Bryant S.V., Stanbridge E.J.

Analysis of the pattern of *QM* expression during mouse development, 64:161-171

Dictyostelium

Hadwiger J.A., Srinivasan J.

Folic acid stimulation of the *G α G* protein-mediated signal transduction pathway inhibits anterior prestalk cell development in *Dictyostelium*, 64:195-204

Agarwal A., Blumberg D.D.

Dictyostelium ribosomal protein genes and the elongation factor 1B gene show coordinate developmental regulation which is under post-transcriptional control, 64:247-254

Differential display

Cram D.S., McIntosh A., Oxbrow L., Johnston A.M.,

DeAizpurua H.

Differential mRNA display analysis of two related but functionally distinct rat insulinoma (RIN) cell lines: identification of CD24 antigen expression in the developing pancreas, 64: 237-246

dlk protein

Garcés C., Ruiz-Hidalgo M.J., Bonvini E., Goldstein J., Laborda J.

Adipocyte differentiation is modulated by secreted delta-like (*dlk*) variants and requires the expression of membrane-associated *dlk*, 64:103-114

Endothelial cells

Zangani D., Darcy K.M., Masso-Welch P.A., Bellamy E.S.,

Desole M.S., Ip M.M.

Multiple differentiation pathways of rat mammary stromal cells *in vitro*: acquisition of a fibroblast, adipocyte or endothelial phenotype is dependent on hormonal and extracellular matrix stimulation, 64:91-101

Epidermis

Compton C.C., Nadire K.B., Regauer S., Simon M., Warland G., Landry D.B.

Cultured human sole-derived keratinocyte grafts re-express site-specific differentiation after transplantation, 64:45-53

Epithelia

Subramanian V., Meyer B., Evans G.S.

The murine *Cdx1* gene product localises to the proliferative compartment in the developing and regenerating intestinal epithelium, 64:11-18

Compton C.C., Nadire K.B., Regauer S., Simon M., Warland G., Landry D.B.

Cultured human sole-derived keratinocyte grafts re-express site-specific differentiation after transplantation, 64:45-53

Kurzrock E.A., Baskin L.S., Cunha G.R.

Ontogeny of the male urethra: Theory of endodermal differentiation, 64:115-122

Frojdman K., Miner J.H., Sanes J.R., Pelliniemi L.J., Virtanen I. Sex-specific localization of laminin $\alpha 5$ chain in the differentiating rat testis and ovary, 64:151-159

Mertens C., Kuhn C., Moll R., Schwetlick I., Franke W.W.
Desmosomal plakophilin 2 as a differentiation marker in normal and malignant tissues, 64:271–290

Schmidt A., Langbein L., Prätzel S., Rode M., Rackwitz H.-R., Franke W.W.
Plakophilin 3 – a novel cell-type-specific desmosomal plaque protein, 64:291–306

Epidermis

Compton C.C., Nadire K.B., Regauer S., Simon M., Warland G., Landry D.B.
Cultured human sole-derived keratinocyte grafts re-express site-specific differentiation after transplantation, 64:45–53

DiSepio D., Bickenbach J.R., Longley M.A., Bundman D.S., Rothnagel J.A., Roop D.R.
Characterization of loricrin regulation in vitro and in transgenic mice, 64:225–235

Mills A.A., Mills M.J., Gardiner D.M., Bryant S.V., Stanbridge E.J.
Analysis of the pattern of *QM* expression during mouse development, 64:161–171

Tomakidi P., Mirancea N., Fusenig N.E., Herold-Mende C., Bosch F.X., Breitkreutz D.
Defects of basement membrane and hemidesmosome structure correlate with malignant phenotype and stromal interactions in HaCaT-Ras xenografts, 64:263–275

Epithelial stromal interactions

Tomakidi P., Mirancea N., Fusenig N.E., Herold-Mende C., Bosch F.X., Breitkreutz D.
Defects of basement membrane and hemidesmosome structure correlate with malignant phenotype and stromal interactions in HaCaT-Ras xenografts, 64:263–275

Epithelial-mesenchymal transition

Muñoz-Chápuli R., Pérez-Pomares J.M., Maciás D., García-Garrido L., Carmona R., González M.
Differentiation of hemangioblasts from embryonic mesothelial cells? A model on the origin of the vertebrate cardiovascular system, 64:133–141

Embryonic stem cells

Thorsteinsdóttir S., Roelen B.A.J., Goumans M.-J., Ward-van Oostwaard D., Gaspar A.C., Mummery C.L.
Expression of the $\alpha 6A$ integrin splice variant in developing mouse embryonic stem cell aggregates and correlation with cardiac muscle differentiation, 64:173–184

Extracellular matrix

Denker A.E., Haas A.R., Nicoll S.B., Tuan R.S.
Chondrogenic differentiation of murine C3H10T1/2 multipotential mesenchymal cells: I. Stimulation by bone morphogenetic protein-2 in high-density micromass cultures, 64:67–76

Zangani D., Darcy K.M., Masso-Welch P.A., Bellamy E.S., Desole M.S., Ip M.M.
Multiple differentiation pathways of rat mammary stromal cells *in vitro*: acquisition of a fibroblast, adipocyte or endothelial phenotype is dependent on hormonal and extracellular matrix stimulation, 64:91–101

Fibroblasts

Zangani D., Darcy K.M., Masso-Welch P.A., Bellamy E.S., Desole M.S., Ip M.M.
Multiple differentiation pathways of rat mammary stromal cells *in vitro*: acquisition of a fibroblast, adipocyte or endothelial phenotype is dependent on hormonal and extracellular matrix stimulation, 64:91–101

Folic acid

Hadwiger J.A., Srinivasan J.
Folic acid stimulation of the $G\alpha 4$ G protein-mediated signal transduction pathway inhibits anterior prestalk cell development in *Dictyostelium*, 64:195–204

G proteins

Hadwiger J.A., Srinivasan J.
Folic acid stimulation of the $G\alpha 4$ G protein-mediated signal transduction pathway inhibits anterior prestalk cell development in *Dictyostelium*, 64:195–204

Gene overexpression

Berberich S.J., Litteral V., Mayo L.D., Tabesh D., Morris D.
mdm-2 gene amplification in 3T3-L1 preadipocytes, 64:205–212

Gonads

Fröjdman K., Miner J.H., Sanes J.R., Pelliniemi L.J., Virtanen I.
Sex-specific localization of laminin $\alpha 5$ chain in the differentiating rat testis and ovary, 64:151–159

Growth factors

Roelen B.A.J., Goumans M.-J., Zwijsen A., Mummery C.L.
Identification of two distinct functions for TGF- β in early mouse development, 64:19–31

Haas A.R., Tuan R.S.
Chondrogenic differentiation of murine C3H10T1/2 multipotential mesenchymal cells: II. Stimulation by bone morphogenetic protein-2 requires modulation of N-cadherin expression and function, 64:77–89

Garcés C., Ruiz-Hidalgo M.J., Bonvini E., Goldstein J., Laborda J.
Adipocyte differentiation is modulated by secreted delta-like (dlk) variants and requires the expression of membrane-associated dlk, 64:103–114

Thorsteinsdóttir S., Roelen B.A.J., Goumans M.-J., Ward-van Oostwaard D., Gaspar A.C., Mummery C.L.
Expression of the $\alpha 6A$ integrin splice variant in developing mouse embryonic stem cell aggregates and correlation with cardiac muscle differentiation, 64:173–184

Hemangioblasts

Muñoz-Chápuli R., Pérez-Pomares J.M., Maciás D., García-Garrido L., Carmona R., González M.
Differentiation of hemangioblasts from embryonic mesothelial cells? A model on the origin of the vertebrate cardiovascular system, 64:133–141

Hematopoiesis

Muñoz-Chápuli R., Pérez-Pomares J.M., Maciás D., García-Garrido L., Carmona R., González M.
Differentiation of hemangioblasts from embryonic mesothelial cells? A model on the origin of the vertebrate cardiovascular system, 64:133–141

Hemidesmosomes

Tomakidi P., Mirancea N., Fusenig N.E., Herold-Mende C., Bosch F.X., Breitkreutz D.
Defects of basement membrane and hemidesmosome structure correlate with malignant phenotype and stromal interactions in HaCaT-Ras xenografts, 64:263–275

Heterokaryon formation

Hirayama E., Nakanishi M., Honda N., Kim J.
Mouse C2 myoblast cells resist HVJ (Sendai virus)-mediated cell fusion in the proliferating stage but become capable of fusion after differentiation, 64:213–223

Immunosurgery

Roelen B.A.J., Goumans M.-J., Zwijsen A., Mummery C.L.
Identification of two distinct functions for TGF- β in early mouse development, 64:19–31

Insulinoma

Cram D.S., McIntosh A., Oxbrow L., Johnston A.M., DeAizpurua H.
Differential mRNA display analysis of two related but functionally distinct rat insulinoma (RIN) cell lines: identification of CD24 antist expression in the developing pancreas, 64:237-246

Integrins

Thorsteinsdóttir S., Roelen B.A.J., Goumans M.-J., Ward-van Oostwaard D., Gaspar A.C., Mummery C.L.
Expression of the $\alpha 6 A$ integrin splice variant in developing mouse embryonic stem cell aggregates and correlation with cardiac muscle differentiation, 64:173-184

Tomakidi P., Mirancea N., Fusenig N.E., Herold-Mende C., Bosch F.X., Breitkreutz D.
Defects of basement membrane and hemidesmosome structure correlate with malignant phenotype and stromal interactions in HaCaT-Ras xenografts, 64:263-275

Intestine

Subramanian V., Meyer B., Evans G.S.
The murine *Cdx1* gene product localises to the proliferative compartment in the developing and regenerating intestinal epithelium, 64:11-18

Islets

Cram D.S., McIntosh A., Oxbrow L., Johnston A.M., DeAizpurua H.
Differential mRNA display analysis of two related but functionally distinct rat insulinoma (RIN) cell lines: identification of CD24 antist expression in the developing pancreas, 64:237-246

Junctions

Mertens C., Kuhn C., Moll R., Schwetlick I., Franke W.W.
Desmosomal plakophilin 2 as a differentiation marker in normal and malignant tissues, 64:271-290

Schmidt A., Langbein L., Prätzel S., Rode M., Rackwitz H.-R., Franke W.W.
Plakophilin 3 - a novel cell-type-specific desmosomal plaque protein, 64:291-306

Keratinocytes

Compton C.C., Nadire K.B., Regauer S., Simon M., Warland G., Landry D.B.
Cultured human sole-derived keratinocyte grafts re-express site-specific differentiation after transplantation, 64:45-53

DiSepio D., Bickenbach J.R., Longley M.A., Bundman D.S., Rothnagel J.A., Roop D.R.
Characterization of loricrin regulation in vitro and in transgenic mice, 64:225-235

Keratins

Tomakidi P., Mirancea N., Fusenig N.E., Herold-Mende C., Bosch F.X., Breitkreutz D.
Defects of basement membrane and hemidesmosome structure correlate with malignant phenotype and stromal interactions in HaCaT-Ras xenografts, 64:263-275

Compton C.C., Nadire K.B., Regauer S., Simon M., Warland G., Landry D.B.
Cultured human sole-derived keratinocyte grafts re-express site-specific differentiation after transplantation, 64:45-53

Zou C.-P., Hong W.K., Lotan R.
Expression of retinoic acid receptor β is associated with inhibition of keratinization in human head and neck squamous carcinoma cells, 64:123-132

McBride S., Walsh D., Meleady P., Daly N., Clynes M.
Bromodeoxyuridine induces keratin protein synthesis at a post-transcriptional level in human lung tumour cell lines, 64:185-193

Laminin

Fröjdman K., Miner J.H., Sanes J.R., Pelliniemi L.J., Virtanen I.
Sex-specific localization of laminin $\alpha 5$ chain in the differentiating rat testis and ovary, 64:151-159

LAZ3/BCL6 Oncogene

Albagli-Curiel O., Dhordain P., Lantoine D., Auradé F., Quief S., Kerckaert J.-P., Montarras D., Pinset C.
Increased expression of the LAZ(BCL6)proto-oncogene accompanies murine skeletal myogenesis, 64:33-44

Lens differentiation

Sullivan C.H., Marker P.C., Thorn J.M., Brown J.D.
Reliability of delta-crystallin as a marker for studies of chick lens induction, 64:1-9

Mizuno N., Mochii M., Takahashi T.C., Eguchi G., Okada T.S.
Lens regeneration in *Xenopus* is not a mere repeat of lens development, with respect to crystallin gene expression, 64:143-149

Lens regeneration

Mizuno N., Mochii M., Takahashi T.C., Eguchi G., Okada T.S.
Lens regeneration in *Xenopus* is not a mere repeat of lens development, with respect to crystallin gene expression, 64:143-149

Liver

Stier H., Fahimi H.D., Van Veldhoven P.P., Mannaerts G.P., Völkl A., Baumgart E.
Maturation of peroxisomes in differentiating human hepatoblastoma cells (HepG2): possible involvement of the peroxisome proliferator-activated receptor α (PPAR α), 64:35-66

Lung

McBride S., Walsh D., Meleady P., Daly N., Clynes M.
Bromodeoxyuridine induces keratin protein synthesis at a post-transcriptional level in human lung tumour cell lines, 64:185-193

Mammary stroma

Zangani D., Darcy K.M., Masso-Welch P.A., Bellamy E.S., Desole M.S., Ip M.M.
Multiple differentiation pathways of rat mammary stromal cells *in vitro*: acquisition of a fibroblast, adipocyte or endothelial phenotype is dependent on hormonal and extracellular matrix stimulation, 64:91-101

mdm-2

Berberich S.J., Litteral V., Mayo L.D., Tabesh D., Morris D.
mdm-2 gene amplification in 3T3-L1 preadipocytes, 64:205-212

Mesothelium

Muñoz-Chápuli R., Pérez-Pomares J.M., Maciás D., García-Garrido L., Carmona R., González M.
Differentiation of hemangioblasts from embryonic mesothelial cells? A model on the origin of the vertebrate cardiovascular system, 64:133-141

Micromass culture

Denker A.E., Haas A.R., Nicoll S.B., Tuan R.S.
Chondrogenic differentiation of murine C3H10T1/2 multipotential mesenchymal cells: I. Stimulation by bone morphogenetic protein-2 in high-density micromass cultures, 64:67-76

Mouse cells and tissues

Subramanian V., Meyer B., Evans G.S.
The murine *Cdx1* gene product localises to the proliferative compartment in the developing and regenerating intestinal epithelium, 64:11-18

Denker A.E., Haas A.R., Nicoll S.B., Tuan R.S.
Chondrogenic differentiation of murine C3H10T1/2 multipotential mesenchymal cells: I. Stimulation by bone morphogenetic protein-2 in high-density micromass cultures, 64:67-76

Haas A.R., Tuan R.S.
Chondrogenic differentiation of murine C3H10T1/2 multipotential mesenchymal cells: II. Stimulation by bone morphogenetic protein-2 requires modulation of N-cadherin expression and function, 64:77-89

Kurzrock E.A., Baskin L.S., Cunha G.R.
Ontogeny of the male urethra: Theory of endodermal differentiation, 64:115-122

Mills A.A., Mills M.J., Gardiner D.M., Bryant S.V., Stanbridge E.J.
Analysis of the pattern of *QM* expression during mouse development, 64:161-171

Thorsteinsdóttir S., Roelen B.A.J., Goumans M.-J., Ward-van Oostwaard D., Gaspar A.C., Mummery C.L.
Expression of the $\alpha 6A$ integrin splice variant in developing mouse embryonic stem cell aggregates and correlation with cardiac muscle differentiation, 64:173-184

Hirayama E., Nakanishi M., Honda N., Kim J.
Mouse C2 myoblast cells resist HVJ (Sendai virus)-mediated cell fusion in the proliferating stage but become capable of fusion after differentiation, 64:213-223

Roelen B.A.J., Goumans M.-J., Zwijsen A., Mummery C.L.
Identification of two distinct functions for TGF- β in early mouse development, 64:19-31

DiSepio D., Bickenbach J.R., Longley M.A., Bundman D.S., Rothnagel J.A., Roop D.R.
Characterization of loricrin regulation in vitro and in transgenic mice, 64:225-235

Myoblast

Hirayama E., Nakanishi M., Honda N., Kim J.
Mouse C2 myoblast cells resist HVJ (Sendai virus)-mediated cell fusion in the proliferating stage but become capable of fusion after differentiation, 64:213-223

Myogenesis

Albagli-Curiel O., Dhordain P., Lantoine D., Auradé F., Quief S., Kerckaert J.-P., Montarras D., Pinset C.
Increased expression of the LAZ(BCL6) proto-oncogene accompanies murine skeletal myogenesis, 64:33-44

Hirayama E., Nakanishi M., Honda N., Kim J.
Mouse C2 myoblast cells resist HVJ (Sendai virus)-mediated cell fusion in the proliferating stage but become capable of fusion after differentiation, 64:213-223

Myotube

Thorsteinsdóttir S., Roelen B.A.J., Goumans M.-J., Ward-van Oostwaard D., Gaspar A.C., Mummery C.L.
Expression of the $\alpha 6A$ integrin splice variant in developing mouse embryonic stem cell aggregates and correlation with cardiac muscle differentiation, 64:173-184

N-cadherin

Haas A.R., Tuan R.S.
Chondrogenic differentiation of murine C3H10T1/2 multipotential mesenchymal cells: II. Stimulation by bone morphogenetic protein-2 requires modulation of N-cadherin expression and function, 64:77-89

Oncogenes

Albagli-Curiel O., Dhordain P., Lantoine D., Auradé F., Quief S., Kerckaert J.-P., Montarras D., Pinset C.
Increased expression of the LAZ(BCL6) proto-oncogene accompanies murine skeletal myogenesis, 64:33-44

Tomakidi P., Mirancea N., Fusenig N.E., Herold-Mende C., Bosch F.X., Breitkreutz D.

Defects of basement membrane and hemidesmosome structure correlate with malignant phenotype and stromal interactions in HaCaT-Ras xenografts, 64:263-275

Mills A.A., Mills M.J., Gardiner D.M., Bryant S.V., Stanbridge E.J.
Analysis of the pattern of *QM* expression during mouse development, 64:161-171

Ovary

Fröjdman K., Miner J.H., Sanes J.R., Pelliniemi L.J., Virtanen I.
Sex-specific localization of laminin $\alpha 5$ chain in the differentiating rat testis and ovary, 64:151-159

p53

Berberich S.J., Litteral V., Mayo L.D., Tabesh D., Morris D.
mdm-2 gene amplification in 3T3-L1 preadipocytes, 64:205-212

Pancreas

Cram D.S., McIntosh A., Oxbrow L., Johnston A.M., DeAizpurua H.
Differential mRNA display analysis of two related but functionally distinct rat insulinoma (RIN) cell lines: identification of CD24 antist expression in the developing pancreas, 64: 237-246

Parietal endoderm

Roelen B.A.J., Goumans M.-J., Zwijsen A., Mummery C.L.
Identification of two distinct functions for TGF- β in early mouse development, 64:19-31

Penis

Kurzrock E.A., Baskin L.S., Cunha G.R.
Ontogeny of the male urethra: Theory of endodermal differentiation, 64:115-122

Peroxisome proliferation

Stier H., Fahimi H.D., Van Veldhoven P.P., Mannaerts G.P., Völkl A., Baumgart E.
Maturation of peroxisomes in differentiating human hepatoblastoma cells (HepG2): possible involvement of the peroxisome proliferator-activated receptor α (PPAR α), 64:35-66

Plakophilins

Mertens C., Kuhn C., Moll R., Schwetlick I., Franke W.W.
Desmosomal plakophilin 2 as a differentiation marker in normal and malignant tissues, 64:271-290

Schmidt A., Langbein L., Prätzel S., Rode M., Rackwitz H.-R., Franke W.W.
Plakophilin 3 - a novel cell-type-specific desmosomal plaque protein, 64:291-306

Plaques

Schmidt A., Langbein L., Prätzel S., Rode M., Rackwitz H.-R., Franke W.W.
Plakophilin 3 - a novel cell-type-specific desmosomal plaque protein, 64:291-306

Mertens C., Kuhn C., Moll R., Schwetlick I., Franke W.W.
Desmosomal plakophilin 2 as a differentiation marker in normal and malignant tissues, 64:271-290

Preadipocytes

Berberich S.J., Litteral V., Mayo L.D., Tabesh D., Morris D.
mdm-2 gene amplification in 3T3-L1 preadipocytes, 64:205-212

Boone C., Grégoire F., Remacle C.
Various stimulators of the cyclic AMP pathway fail to promote adipose conversion of porcine preadipocytes in primary culture, 64:255-262

Garcés C., Ruiz-Hidalgo M.J., Bonvini E., Goldstein J., Laborda J. Adipocyte differentiation is modulated by secreted delta-like (dlk) variants and requires the expression of membrane-associated dlk, 64:103-114

QM expression

Mills A.A., Mills M.J., Gardiner D.M., Bryant S.V., Stanbridge E.J. Analysis of the pattern of QM expression during mouse development, 64:161-171

Rat cells and tissues

Zangani D., Darcy K.M., Masso-Welch P.A., Bellamy E.S., Desole M.S., Ip M.M. Multiple differentiation pathways of rat mammary stromal cells *in vitro*: acquisition of a fibroblast, adipocyte or endothelial phenotype is dependent on hormonal and extracellular matrix stimulation, 64:91-101

Kurzrock E.A., Baskin L.S., Cunha G.R. Ontogeny of the male urethra: Theory of endodermal differentiation, 64:115-122

Fröjdman K., Miner J.H., Sanes J.R., Pelliniemi L.J., Virtanen I. Sex-specific localization of laminin $\alpha 5$ chain in the differentiating rat testis and ovary, 64:151-159

Cram D.S., McIntosh A., Oxbrow L., Johnston A.M., DeAizpurua H. Differential mRNA display analysis of two related but functionally distinct rat insulinoma (RIN) cell lines: identification of CD24 antist expression in the developing pancreas, 64: 237-246

Receptors

Roelen B.A.J., Goumans M.-J., Zwijsen A., Mummery C.L. identification of two distinct functions for TGF- β in early mouse development, 64:19-31

Stier H., Fahimi H.D., Van Veldhoven P.P., Mannaerts G.P., Völkl A., Baumgart E. Maturation of peroxisomes in differentiating human hepatoblastoma cells (HepG2): possible involvement of the peroxisome proliferator-activated receptor α (PPAR α), 64:35-66

Zou C.-P., Hong W.K., Lotan R. Expression or retinoic acid receptor β is associated with inhibition of keratinization in human head and neck squamous carcinoma cells, 64:123-132

Retinoic acid

Zou C.-P., Hong W.K., Lotan R. Expression or retinoic acid receptor β is associated with inhibition of keratinization in human head and neck squamous carcinoma cells, 64:123-132

Retinoic acid receptor β

Zou C.-P., Hong W.K., Lotan R. Expression or retinoic acid receptor β is associated with inhibition of keratinization in human head and neck squamous carcinoma cells, 64:123-132

Ribosomes

Agarwal A., Blumberg D.D. *Dictyostelium* ribosomal protein genes and the elongation factor 1B gene show coordinate developmental regulation which is under post-transcriptional control, 64:247-254

Mills A.A., Mills M.J., Gardiner D.M., Bryant S.V., Stanbridge E.J. Analysis of the pattern of QM expression during mouse development, 64:161-171

RT-PCR

Thorsteinsdóttir S., Roelen B.A.J., Goumans M.-J., Ward-van Oostwaard D., Gaspar A.C., Mummery C.L. Expression of the $\alpha 6A$ integrin splice variant in developing mouse embryonic stem cell aggregates and correlation with cardiac muscle differentiation, 64:173-184

Sex differentiation

Fröjdman K., Miner J.H., Sanes J.R., Pelliniemi L.J., Virtanen I. Sex-specific localization of laminin $\alpha 5$ chain in the differentiating rat testis and ovary, 64:151-159

Skin

Compton C.C., Nadire K.B., Regauer S., Simon M., Warland G., Landry D.B. Cultured human sole-derived keratinocyte grafts re-express site-specific differentiation after transplantation, 64:45-53

Smad

Roelen B.A.J., Goumans M.-J., Zwijsen A., Mummery C.L. identification of two distinct functions for TGF- β in early mouse development, 64:19-31

Squamous cell carcinoma

Zou C.-P., Hong W.K., Lotan R. Expression or retinoic acid receptor β is associated with inhibition of keratinization in human head and neck squamous carcinoma cells, 64:123-132

Testis

Fröjdman K., Miner J.H., Sanes J.R., Pelliniemi L.J., Virtanen I. Sex-specific localization of laminin $\alpha 5$ chain in the differentiating rat testis and ovary, 64:151-159

Transcription rate analysis

Agarwal A., Blumberg D.D. *Dictyostelium* ribosomal protein genes and the elongation factor 1B gene show coordinate developmental regulation which is under post-transcriptional control, 64:247-254

Transcriptional repressor

Albagli-Curiel O., Dhordain P., Lantoine D., Auradé F., Quief S., Kerckaert J.-P., Montarras D., Pinset C. Increased expression of the LAZ(BCL6)proto-oncogene accompanies murine skeletal myogenesis, 64:33-44

Transforming growth factor β

Thorsteinsdóttir S., Roelen B.A.J., Goumans M.-J., Ward-van Oostwaard D., Gaspar A.C., Mummery C.L. Expression of the $\alpha 6A$ integrin splice variant in developing mouse embryonic stem cell aggregates and correlation with cardiac muscle differentiation, 64:173-184

Haas A.R., Tuan R.S.

Chondrogenic differentiation of murine C3H10T1/2 multipotential mesenchymal cells: II. Stimulation by bone morphogenetic protein-2 requires modulation of N-cadherin expression and function, 64:77-89

Denker A.E., Haas A.R., Nicoll S.B., Tuan R.S.

Chondrogenic differentiation of murine C3H10T1/2 multipotential mesenchymal cells: I. Stimulation by bone morphogenetic protein-2 in high-density micromass cultures, 64:67-76

Transgenic animals

DiSepio D., Bickenbach J.R., Longley M.A., Bundman D.S., Rothnagel J.A., Roop D.R. Characterization of loricrin regulation *in vitro* and in transgenic mice, 64:225-235

Tumor invasion

Tomakidi P., Mirancea N., Fusenig N.E., Herold-Mende C., Bosch F.X., Breitkreutz D.

Defects of basement membrane and hemidesmosome structure correlate with malignant phenotype and stromal interactions in HaCaT-Ras xenografts, 64:263-275

Two-cell block

Roelen B.A.J., Goumans M.-J., Zwijsen A., Mummery C.L. identification of two distinct functions for TGF- β in early mouse development, 64:19-31

Urethra

Kurzrock E.A., Baskin L.S., Cunha G.R.

Ontogeny of the male urethra: Theory of endodermal differentiation, 64:115-122

Vasculogenesis

Zangani D., Darcy K.M., Masso-Welch P.A., Bellamy E.S., Desole M.S., Ip M.M.

Multiple differentiation pathways of rat mammary stromal cells *in vitro*: acquisition of a fibroblast, adipocyte or endothelial phenotype is dependent on hormonal and extracellular matrix stimulation, 64:91-101

Muños-Chápuli R., Pérez-Pomares J.M., Maciás D., García-Garrido L., Carmona R., González M.

Differentiation of hemangioblasts from embryonic mesothelial cells? A model on the origin of the vertebrate cardiovascular system, 64:133-141

Xenopus

Mizuno N., Mochii M., Takahashi T.C., Eguchi G., Okada T.S.

Lens regeneration in *Xenopus* is not a mere repeat of lens development, with respect to crystallin gene expression, 64:143-149